

International Journal of Current Microbiology and Applied Sciences ISSN: 2319-7706 Volume 8 Number 09 (2019)

Journal homepage: http://www.ijcmas.com



Original Research Article

https://doi.org/10.20546/ijcmas.2019.809.130

Assess the Performance of Different Breeds under Backyard Poultry Farming System in Erode District, India

S. Vinothraj¹*, P. Alagesan² and M. Siva²

¹Department of Animal Science, ²Department of Home Science, ICAR- Krishi Vigyan Kendra (MYRADA), Erode District, TamilNadu-638453, India

*Corresponding author

ABSTRACT

Keywords

Body weight, Desi chickens, Egg production, Mortality, Gramapriya, Srinidhi

Article Info

Accepted: 20 August 2019 Available Online: 10 September 2019 A study was conducted to assess the Performance of Different Breeds which includes Gramapriya, TANUVAS Aseel and Srinidhi under Backyard Poultry farming System in Erode district by covering 30 backyard poultry farmers in T.N.Palayam, Gobi and Anthiyur block of Erode district, Tamil Nadu. Information was obtained on mean body weight at various ages, age at first egg, annual egg production and mortality rate. The result indicated that Gramapriya showed better annual egg production and Srinidhi gained better body weight at 20 th week when compared to other strains. Even though the two strains performed well in productivity, TANUVAS Aseel is suitable for backyard farming system due its better farmers adoption due its external appearance and genetic potentiality. Attempts were taken in 2018-19 to link the results of the strain assessment to the mainstream extension at the district for larger adoption of rural communities.

Introduction

Backyard poultry production is an old age profession of rural families of India. It is the most potent source for subsidiary incomes for landless and poor farmers. It is an enterprise with low initial investment but higher economic returns and can easily be managed by women, children and old aged persons of the households.

Now-a-days, poultry meat and eggs have been the best and cheapest sources for meeting out the per capita requirement of protein and

energy for rural areas of India. Though India has shown a tremendous growth in poultry production over decades but rural poultry farming is still lagging behind and always found neglected. As it is the best alternative the small scale farmers subsidies the income with negligible input, this farming system needs an upliftment with newly developed varieties of chicken. Rural farmers usually rear desi type chicken having low egg and meat production potential. Most of the backyard poultry production comprises of rearing indigenous birds with poor production performances (Pathak and Nath, 2013; Chakravarthi et al.,., 2014; Reetha et al.,, 2016; Patra and Singh, 2016). However, over the period of time improved strains have introduced been by extension agencies. development Development organizations under government of India have developed improved strains like Gramapriya, TANUVAS Aseel and Srinidhi.

Gramapriya is a multicolored egg purpose chicken variety developed at Directorate of Poultry Research, Hyderabad for free range and rural backyard rearing. TANUVAS Aseel is a new variety of native chicken developed at Poultry Research Station, TANUVAS. Srinidhi is a dual purpose chicken variety developed at Directorate of Poultry Research, Hyderabad.

By understanding the potentiality of the above strains KVK conducted an OFT in 2018-2019 to assess its suitability and performance in the backyard farming system to feed into the mainstream extension. This paper is based an results of the OFT conducted by ICAR, KVK MYRADA with the following objectives.

To study body weight at different ages. To find out the age at egg laying of this particular study. To assess the annual egg production . To study the mortality rate and also to identify the farmers adoptability.

Materials and Methods

The present study was conducted in the Erode district situated between 10-35' and 11-60' of north latitude and 76.49' and 77.58' of East longitude and 171-91' meters above the mean sea level. The river Cauvery flows on the north and eastern part of the District. Erode town sweats under very hot spells during summer. The study was carried out in Gobi, T.N. Palayam and Anthiyur blocks of Erode district of Tamil Nadu during 2018-2019.

The selected Poultry farmers were trained on all scientific Desi bird training techniques like brooding, deworming, Vacciantion etc. Each farmer supported with of 30 unsexed day old chicks which includes 10 Gramapriya chicks, 10 TANUVAS Aseel Chicks and 10 Srinidhi Chicks. There improved strains compared compared with farmers traditional breeds in this study.

The KVK Scientist made regular visit to farmer's field and observed production parameters.

The following production parameters were studied

Mortality percentage 0-12 weeks

Body weight at 12 th week age (Kg)

Body weight at 20 th week age (Kg)

Average annual egg production

Average age at first egg

Simple percentage analysis was used to analyze the data.

Results and Discussion

The production parameters of these improved

strains of chicken are presented in Table.1

Mortality Percentage (0-12 Weeks)

Higher mortality rate were recorded in improved strain than desi chicks during the early part of the life (0-12 Weeks). The reason for lower mortality in desi chicks might be due to more adaptability to the environment and proper brooding by their mother.

Body weight at 20 th week

The body weight at 20 th week is a factor which ensure the market value of the birds in the present study. Srinidhi attained better body weight & 20 th week (1.6 Kg) when compared to other two strains. The present findings were in accordance with findings of Sharma *et al.*,, (2017) who reported that 20 th week body weight of Srindhi birds under backyard system of poultry rearing was 1.68 Kg.

Annual Average Egg production

Egg production determining the success of Poultry enterprises and gaining better income to farmers. Comparative estimate of egg production revealed that Gramapriya has higher annual egg production (182 eggs) when compared to other two strains. The variation in egg production might be due to genetic potential of Native and improved varieties of Chicken. Karuna sree *et al.*, (2017), who reported egg production (178 eggs / year) of Gramapriya, has similar results of the present study.

Age at First egg

TANUVAS Aseel started to lay egg at 155 days of age but Gramapriya and Srinidhi started to lay eggs at 162 days and 188 days respectively.

Table.1Comparative performance of Gramapriya, TANUVAS Aseel and Srinidhi under backyard system of rearing

S. No	Particulars	Mortality percentage (0-12 Weeks)	Body weight at 12 th week	Body weight at 20 th week	Average age at egg laying (Days)	Annual average egg production
1.	Farmers practice	2	0.89 Kg	1.24 Kg	194	70
2.	Gramapriya	5.8	1.11 Kg	1.54 Kg	162	182
3.	TANUVAS Aseel	8.2	1.02 Kg	1.38 Kg	155	151
4.	Srinidhi	7.9	1.1 Kg	1.60 Kg	188	150

From the study, it can be concluded that Gramapriya and Srinidhi birds performs better than TANUVAS Aseel chicken in terms of annual egg production and body weight under backyard system of rearing, But TANUVAS Aseel breed had better market opportunity and farmers adoption due its external appearance and genetic potentiality. So, farmers from rural areas of Erode district can

rear TANUVAS Aseel for their livelihood and nutritional security.

References

Pathak PK and Nath BG. Rural Poultry Farming with Improved Breed of Backyard Chicken. *Journal of* World"s Poultry Research, 2013; 3

- (1): 24-273.
- Chakravarthy PV, Mohan B and Senthilkumar K. Performance of CARI Nirbheek (Aseel Cross) birds reared under intensive and semi intensive system in Namakkal District. *Indian Veterinary Journal*, 2014; 91(11): 85-87.
- Patra, J and Singh DV (2016). Backyard poultry farming, a suitable Intervention for Tribal people for their livelihood support and Nutritional security. International Journal of Humanities and Social Science Innovation, 2016; 5(6): 22-26
- Reetha TL, Rajeswar JJ, Harikrishnan TJ, Sukumar K, Srinivasan P, Kirubakaran JJ (2016) Studies on the effectiveness of oral pellet vaccine in improving egg production and egg quality in desi chicken. *Veterinary World*, 2016; 9 (8): 900-903.

- Karuna sree E, Vijaya Nirmala T and Reddy RVSK. Comparative performance of different varieties of chicken under backyard system of rearing in tribal and rural areas of West Godavari district, A.P. *International Journal of Science, Enviro nment and Technology*, 2017; 6(2): 1100 1104
- Sarma M, Islam R, Borah MK, Sharma P, Mahanta JD. Kalita N BN. Bhattacharyya Comparative performance of Vanaraja, Srinidhi and Desi chicken under traditional system among tribal community of Assam Comparative performance of Vanaraja, Srinidhi and Desi chicken under traditional system among tribal community of Assam. Indian Journal of Animal Research, 2017; B-3391:1-3

How to cite this article:

Vinothraj, S., P. Alagesan and Siva, M.. 2019. Assess the Performance of Different Breeds under Backyard Poultry Farming System in Erode District. *Int.J.Curr.Microbiol.App.Sci.* 8(09): 1138-1141. doi: https://doi.org/10.20546/ijcmas.2019.809.130